

## **End-tidal CO<sub>2</sub> pressure: an important parameter for a correct interpretation of changes in cerebral hemodynamics and oxygenation measured with functional near infrared spectrophotometry (fNIRS)**

Scholkmann F<sup>1,2</sup>, Gerber U<sup>1,3</sup>, Klein SD<sup>1</sup>, Wolf U<sup>1</sup>.

<sup>1</sup>Institute of Complementary Medicine KIKOM, University of Bern, Switzerland

<sup>2</sup>Biomedical Optics Research Laboratory, University Hospital Zurich, Switzerland

<sup>3</sup>University of Lausanne, Switzerland

The aim of the present study was to investigate the effects of different speech tasks (recitation of prose (PR), alliteration (AR) and hexameter (HR) verses) and a control task (mental arithmetic (MA) with voicing of the result) on end-tidal CO<sub>2</sub> (ET-CO<sub>2</sub>), cerebral hemodynamics; i.e. total hemoglobin (tHb) and tissue oxygen saturation (StO<sub>2</sub>). tHb and StO<sub>2</sub> were measured with a frequency domain near infrared spectrophotometer (ISS Inc., USA) and ET-CO<sub>2</sub> with a gas analyzer (Nellcor N1000). Measurements were performed in 24 adult volunteers (11 female, 13 male; age range 22 to 64 years) during task performance in a randomized order on 4 different days to avoid potential carry over effects. Statistical analysis was applied to test differences between baseline, 2 recitation and 5 recovery periods. The two brain hemispheres and 4 tasks were tested separately. Data analysis revealed that during the recitation tasks (PR, AR and HR) StO<sub>2</sub> decreased statistically significant ( $p < 0.05$ ) during PR and AR in the right prefrontal cortex (PFC) and during AR and HR in the left PFC. tHb showed a significant decrease during HR in the right PFC and during PR, AR and HR in the left PFC. During the MA task, StO<sub>2</sub> increased significantly. A significant decrease in ET-CO<sub>2</sub> was found during all 4 tasks with the smallest decrease during the MA task. In conclusion, we hypothesize that the observed changes in tHb and StO<sub>2</sub> are mainly caused by an altered breathing during the tasks that led a lowering of the CO<sub>2</sub> content in the blood provoked a cerebral CO<sub>2</sub> reaction, i.e. a vasoconstriction of blood vessels due to decreased CO<sub>2</sub> pressure and thereby decrease in cerebral blood volume. Therefore, breathing changes should be monitored during brain studies involving speech when using functional near infrared spectroscopy (fNIRS) to ensure a correct interpretation of changes in hemodynamics and oxygenation.